

FINAL

**Appendix B
(Nutrient Limitation Analysis)**

**Total Maximum Daily Loads of Nitrogen for
Three Tidal Tributaries and
Total Maximum Daily Load of Biochemical Oxygen Demand for
One Tributary in the Newport Bay System
Worcester County, Maryland**

FINAL

December 2002

FINAL

Nitrogen and phosphorus are essential nutrients for algal growth. However, common types of algae require different amounts of these two nutrients. If one nutrient is available in great abundance relative to the other nutrient, then the nutrient that is less available restricts the amount of plant that can be produced, regardless of the amount of the other nutrient that is available. This less available nutrient is called the “limiting nutrient”. Before conducting any analysis in terms of reduction, it is necessary to examine the ratio of nitrogen to phosphorus to establish which nutrient is most likely limiting the algal growth in the system.

In general, an DIN:DIP ratio in the range of 7:1 to 12:1 by mass is associated with plant growth being limited by neither phosphorus nor nitrogen. If the N:P ratio is greater than 12:1, phosphorus tends to be limiting, and if the N:P ratio is less than 7:1, nitrogen tends to be limiting (Wetzel, 2001). However only the DIN:DIP ratio by itself is not sufficient to establish which nutrient will be limiting. The limiting nutrient also depends on the concentration of the nutrients as well as bio-availability of these nutrients. Detailed modeling sensitivity runs are also required to justify which nutrient is most likely limiting under existing and other conditions.

Various field data (MDE, DNR, MCBP, NPKS) were examined to analyze the nutrient limitation in the Newport Bay System. The result was analyzed using a ratio of dissolved nitrogen to dissolved phosphorus concentrations. A detailed analysis is presented in the Table B1 below for both low summer sampling and high winter sampling flow conditions. The table suggests that the low flow system is more likely nitrogen limiting even though sometimes the data show it could be phosphorus limiting. The results from the high flow monitoring data suggest the system is kind of borderline, and it is difficult to draw conclusion as to which nutrient will be limiting. A set of sensitivity tests using the WASP model were also conducted using the critical loads and the flows. The results are presented in Table B2.

Table B1: Nutrient Limitation Analysis from Field Monitoring Data

Summer Sampling Data			Winter Sampling Data		
Date	Station	DIN/DIP	Date	Station	DIN/DIP
07/01/1998	AYR0017	2	04/29/1998	AYR0017	3
07/28/1998	AYR0017	1	04/29/1998	AYR0033	132
08/26/1998	AYR0017	0	04/29/1998	KIT0015	30
09/30/1998	AYR0017	4	04/29/1998	NPC0012	19
10/21/1998	AYR0017	4	04/29/1998	TRC0043	11
07/28/1998	AYR0033	2	04/29/1998	XCM1562	4
08/26/1998	AYR0033	3	04/29/1998	TRC0059	59
09/30/1998	AYR0033	3	04/29/1998	XCM1562	4
10/21/1998	AYR0033	30			
07/01/1998	KIT0015	20			
07/28/1998	KIT0015	3			
08/26/1998	KIT0015	26			
09/30/1998	KIT0015	7			
10/21/1998	KIT0015	4			

Summer Sampling Data			Winter Sampling Data		
Date	Station	DIN/DIP	Date	Station	DIN/DIP
07/28/1998	NPC0012	1			
08/26/1998	NPC0012	1			
09/30/1998	NPC0012	2			
07/28/1998	NPC0031	7			
08/26/1998	NPC0031	1			
09/30/1998	NPC0031	2			
10/21/1998	NPC0031	16			
07/01/1998	TRC0043	2			
07/28/1998	TRC0043	0			
08/26/1998	TRC0043	0			
09/30/1998	TRC0043	1			
10/21/1998	TRC0043	1			
08/26/1998	TRC0059	16			
09/30/1998	TRC0059	13			
10/21/1998	TRC0059	4			
07/01/1998	XCM1562	1			
07/28/1998	XCM1562	4			
08/26/1998	XCM1562	0			
09/30/1998	XCM1562	6			
10/21/1998	XCM1562	8			
07/01/1998	XCM3367	3			
07/28/1998	XCM3367	11			
08/26/1998	XCM3367	4			
09/30/1998	XCM3367	5			
10/21/1998	XCM3367	4			
07/01/1998	XCM4878	5			
07/28/1998	XCM4878	5			
08/26/1998	XCM4878	3			
09/30/1998	XCM4878	2			
10/21/1998	XCM4878	4			

The Table B1 above suggests that the system is not likely nitrogen limited in the summer, but it can be either (N or P) limiting in the spring and in the winter. However DIN/DIP ratio itself cannot be sufficient to interpret which is the limiting nutrient. Keeping this in mind, the model was run for different loads of nitrogen and phosphorus for each condition of flow. Initially, the model was run for current loading conditions. Then they were re-run with a double load of phosphorus and a double load of nitrogen separately to see which nutrient affects the system. Similar runs were also conducted to check the impact there would be of cutting the load of individual nutrients (nitrogen or phosphorus) by 50%. Figure B1 and

FINAL

Figure B2 below presents the impact of doubling the nitrogen load and the phosphorus load individually. The run shows that doubling of nitrogen load changed the algal concentration significantly while doubling phosphorus load had a very little effect. Similar results were obtained when the individual loads were reduced by 50% (Figure B3 and Figure B4). This supports the assumption that the system is nitrogen limited during a low flow condition.

Table B2: Nutrient Limitation Analysis from the Model Results

WQ Segments	DIN/DIP		
	Summer	Spring	Winter
1	1	1	3
2	1	2	5
3	2	3	7
4	2	4	9
5	1	7	10
6	1	9	10
7	1	10	10
8	1	11	10
9	1	10	10
10	0	10	11
11	0	11	11
12	7	14	15
20	1	9	9
21	1	7	8
22	1	6	7
25	2	5	9
26	1	1	9
27	1	1	9

Newport Bay
Summer Flow Sensitivity Runs - Nitrogen Load Doubled




Figure B1: Sensitivity Test of Doubling the Nitrogen Load in the Low Flow Condition

Newport Bay
Summer Flow Sensitivity Runs - P Loads Doubled




Figure B2: Sensitivity Test of Doubling the Phosphorus Load in the Low Flow Condition

Newport Bay
Summer Flow Sensitivity Runs - Nitrogen Load Reduced by 50%




Figure B3: Sensitivity Test of Halving the Nitrogen Load in the Low Flow Condition

Newport Bay
Summer Flow Sensitivity Runs - P Loads Reduced by 50%




Figure B4: Sensitivity Test of Halving the Phosphorus Load in the Low Flow Condition

FINAL

Similar runs were conducted for individual nutrients (N or P) for annual average condition of flow to see what would limit the algal growth in the system. Doubling and reducing the nitrogen load by 50% made a significant difference (Figure B5 & Figure B7), while the results were not significant when doubling or reducing to half the phosphorus loads (Figure B6, Figure B8).

Newport Bay
Average Annual Flow Sensitivity Runs - Nitrogen Load Doubled




Figure B5: Sensitivity Test of Doubling the Nitrogen Load in the Average Annual Flow Condition

Newport Bay
Average Annual Flow Sensitivity Runs - Phosphorus Load Doubled




Figure B6: Sensitivity Test of Doubling the Phosphorus Load in the Average Annual Flow Condition

Newport Bay
Average Annual Flow Sensitivity Runs - Nitrogen Load Reduced by 50%




Figure B7: Sensitivity Test of Halving the Nitrogen Load in the Average Annual Flow Condition

Newport Bay
Average Annual Flow Sensitivity Runs - Phosphorus Load Reduced by 50%




Figure B8: Sensitivity Test of Halving the Phosphorus Load in the Average Annual Flow Condition

FINAL

References:

Ambrose, Robert B., Tim A. Wool, James A. Martin. "The Water Quality Analysis Simulation Program, Wasp5". Environmental Research Laboratory, Office Of Research And Development, U.S. Environmental Protection Agency. 1993.

Cerco, C. F., B. Bunch, M. A. Cialone, H. Wang, "Hydrodynamic and Eutrophication Model Study of Indian River and Rehoboth Bay, Delaware," U. S. Army Corps of Engineers Waterways Experimental Station, Philadelphia, PA, Technical Report EL-94-5, May 1994.

Cerco, Carl F. *Water Quality in a Virginia Potomac Embayment: Gunston Cove*. College of William and Mary, Virginia Institute of Marine Science, Gloucester Point, Virginia. April 1985.

Clark L. J., and S. E. Roesh, *Assessment of 1977 Water Quality Conditions in the Upper Potomac Estuary*. U.S. EPA Annapolis Field Office, Annapolis Maryland. EPA 903/9-78-008, 1978.

Coastal Bays Volunteer Monitoring Program, Water Quality Monitoring Manual Maryland Coastal Bays Program, 1997

Di Toro, D.M., J.J. Fitzpatrick, and R.V. Thomann. *Documentation for Water Quality Analysis Simulation Program (WASP) and Model Verification Program (MVP)*. EPA/600/3-81-044. 1983.

Dillow, J., A. Greene, "Ground-Water Discharge and Nitrate Loadings to the Coastal Bays of Maryland" USGS Water Resource Investigations Report 99-4167, 1999.

Dillow, J., A. Greene , U.S.G.S., personal communications, June 2002.

Domotor, Diana K., Michael S. Haire, Narendra N. Panday, and Harry V. Wang. *Mattawoman Creek Water Quality Model*. Technical Report No. 64, Maryland Department of the Environment, Water Management Administration, Modeling and Analysis Division. October 1987.

Haire, M. S., and N. N. Panday, "Quality Assurance/ Quality Control Plan: Water Quality Assessment of the Mattawoman Creek and nearby Potomac Estuary," Office of Environmental Programs, State of Maryland, April 1985.

Institute of Natural Resources, "Sediment Oxygen Demand - Processes, Modeling & Measurement," University of Georgia, Athens, GA, 1986.

Maryland Department of the Environment, Point Source Database, February 2002.

FINAL

Maryland Department of the Environment, "TMDL Quality Assurance Project Plan: Eutrophication Sampling Component," Field Operations Program, Water Quality Monitoring Division, Annapolis Maryland, April 2001.

Maryland Geological Survey, personal communications with Darlene Wells, March 2002.

Mirsajada, Hassan, results of Delaware Inland Bays SOD Survey in 1995 and 1996, August 2000.

National Park Service, personal communication, Brian Sturgis, March 2002

Panday, Narendra N., and Michael S. Haire. *Water Quality Assessment of Mattawoman Creek and the Adjacent Potomac River: Summer 1985*. Technical Report No. 52, Water Management Administration, Modeling and Analysis Division, Maryland Office of Programs, Department of Health and Mental Hygiene. September 1986.

Sampou, Pete correspondence to Friends of Herring and Turville Creeks presenting results of June 5, 1994 Sediment Oxygen Demand Sampling.

Seitzinger, S. P., R. DeKorsey, "Sediment-Water Nutrient Interactions in Rehoboth and Indian River Bays," Academy of Natural Sciences Division of Environmental Research, Philadelphia, PA, submitted to the Inland Bays Estuary Program, Dover DE, June 1994.

Thomann, Robert V., John A. Mueller "Principles of Surface Water Quality Modeling and Control," HarperCollins Publisher Inc., New York, 1987.

Thomann R. V., and J. J. Fitzpatrick. *Calibration and Verification of a Mathematical Model of the Eutrophication of the Potomac Estuary*. HydroQual, Inc. Final Report Prepared for the D.C. Department of Environmental Services, 1982.

Tyson Foods Inc., Berlin– 001 Nitrite, Nitrate Data for 1998, correspondence from John Lister May, 2001.

University of Maryland System, Center for Environmental and Estuarine Studies, "Maryland's Coastal Bays: An assessment of Aquatic Ecosystems, Pollutant Loadings, and Management Options," Chesapeake Biological Laboratory, Solomons, MD, March 1993.

University of Maryland System, Center for Environmental and Estuarine Studies, "Pocomoke River and Sound, Manokin and Big Annemessex Rivers TMDL Project 1999: Water Quality Monitoring Program," Chesapeake Biological Laboratory, Solomons, MD, December 1999.

U.S. EPA, "Technical Guidance Manual for Developing Total Maximum Daily Loads, Book2: Streams and Rivers, Part 1: Biochemical Oxygen Demand/ Dissolved Oxygen and Nutrients/ Eutrophication," Office of Water, Washington D.C., March 1997.

FINAL

U.S. EPA Chesapeake Bay Program. *Chesapeake Bay Program: Watershed Model Application to Calculate Bay Nutrient Loadings: Final Findings and Recommendations*. and Appendices, May, 1996.

Wetzel, R. G., "Limnology: Lake and River Ecosystems," Academic Press, New York, 2001.